




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 7/00	A2	(11) International Publication Number: WO 00/13654 (43) International Publication Date: 16 March 2000 (16.03.00)
<p>(21) International Application Number: PCT/IL99/00478</p> <p>(22) International Filing Date: 5 September 1999 (05.09.99)</p> <p>(30) Priority Data: 126076 4 September 1998 (04.09.98) IL</p> <p>(71) Applicant (for all designated States except US): I.B.R. ISRAELI BIOTECHNOLOGY RESEARCH, LTD. [IL/IL]; 85 Derech Namir, Apt. 23, 62301 Tel Aviv (IL).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SOUDANT, Etienne [FR/FR]; 14, rue Leon Bernard, F-94260 Fresnes (FR). BEZALEL, Lea [IL/IL]; Derech Mezada 15, 84359 Beer Sheva (IL). SCHICKLER, Hedva [IL/IL]; 19 Hamagen Street, 76804 Mazkeret Batya (IL). PALTIEL, Judith [IL/IL]; 12/26 Parshani Street, 76000 Rehovot (IL). BEN-AMOTZ, Ami [IL/IL]; 21 Hatomer Street, 56542 Savion (IL). SHAISH, Aviv [IL/IL]; 37 Achva Street, 79800 Shikmim (IL). PERRY, Inon [IL/IL]; 85 Derech Namir, Apt 23, 62301 Tel Aviv (IL).</p> <p>(74) Agent: REINHOLD COHN AND PARTNERS; P.O.Box 4060, 61040 Tel Aviv (IL).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>

(54) Title: A CAROTENOID PREPARATION

(57) Abstract

A preparation comprising the carotenoids phytoene and phytofluene in an amount which, in combination, is effective in preventing various kinds of damage resulting from oxidation and exposure to UV light is provided. The preparation is essentially colorless and may be used as a topical cosmetic or pharmaceutical preparation as well as an additive in food preparations. In addition, a method for producing substantive amounts of phytoene and phytofluene from carotenoid producing organisms is described as well.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

A CAROTENOID PREPARATION

FIELD OF THE INVENTION

The present invention relates to a carotenoid comprising preparation. The preparation of the invention can be used in food, in cosmetics, in a variety of medicinal uses, etc. The invention also concerns a method for the production of the
5 carotenoids.

BACKGROUND OF THE INVENTION

Carotenoids are pigments produced by microorganisms, fungi and plants, and used by them as antioxidants and protectants against excessive radiation. The most widely used carotenoids in food, medical preparations or cosmetics are
10 β -carotene and lycopene. β -Carotene and lycopene are sensitive to light and oxidation, a property which considerably limits their use and shortens shelf-life of products containing them (in: Carotenoids, Chemistry and Biology, Krinski, N.I., Matthews-Roth, M.M., Taylor, R.F., (Eds), Plenum Press, New York, London, 1989). In addition, β -carotene and lycopene have a distinctive orange color and this
15 color has a serious limitation for a variety of cosmetic or food applications.

Phytoene (7, 8, 11, 12, 7', 8', 11', 12'-octahydro- γ,γ -carotene) and phytofluene (15Z, 7, 8, 11, 12, 7', 8'-hexahydro- γ,γ -carotene), are carotenoids (C-40 isoprenoids chain) which are precursors in the biosynthetic pathway which leads to the production of β -carotene, lycopene and other carotenoids (phytoene is the first

carotenoid-specific precursor and phytofluene is produced therefrom in a subsequent desaturation step). Phytoene is completely colorless whereas phytofluene has a slight yellowish color. Japanese Patent Application No. 90-40520, disclosed that the introduction of a DNA sequence yielding expression of phytoene into certain transfected cancer cells, resulted in inhibition of their growth and in inhibition in the activation of Epstein Barr virus (EBV).

SUMMARY OF THE INVENTION

In accordance with the present invention it was shown that phytoene and phytofluene possess anti-oxidative properties and furthermore are capable of absorbing ultra violet (UV) light. In addition, although having these properties, phytoene and phytofluene were found to be much more stable to oxidation than, for example, β -carotene. These findings led to the realization that phytoene and phytofluene in combination could be useful in the prevention of environmentally induced damage of various kinds.

Thus, in accordance with the invention there is provided a composition comprising, an amount of phytoene and an amount of phytofluene, effective in combination in the prevention of damage resulting from oxidation and from exposure to UV light.

In accordance with the invention, the term "damage" is to be understood as any damage resulting from a variety of oxidative agents such as oxygen itself, hydroxyl radical, hydrogen peroxide, other free radicals, ozone etc., or from any kind of harmful UV irradiation, such as UVA and UVB irradiation. The damage would depend upon the target for which the preparation is used. Thus, if the preparation is used on skin, damage may be any skin damage such as burns, blisters, damage appearing after chronic exposure to sun, e.g. premature aging of the skin, etc. When the preparation is used as a food preservative, such damage may be in the form of a decrease in product stability, chemical modification resulting for example in rancidity, accelerated aging, etc.

In accordance with a preferred embodiment of the invention there is provided a topical skin composition for protecting the skin against environmental hazards, comprising phytoene and phytofluene in an effective amount such that, in combination, these carotenoids exert an oxidation-protecting and UV-protecting effect on the skin.

The term "*environmental hazards*" relates to any environmental agent which can exert damage such as UV radiation or oxidative agents.

The term "*effective amount*" should be understood to mean an amount of phytoene and an amount of phytofluene which, when administered in combination, achieves the desired protective effect.

The phytoene and phytofluene in the composition of the invention may each be either in their *trans* or in their *cis* forms.

The weight ratio between the phytoene and phytofluene in the composition of the invention can range between about 200:1 to about 1:200, respectively, typically between about 50:1 to about 1:50, about 10:1 (phytoene:phytofluene) being a particular example. The above ratios of phytoene to phytofluene may be reached either by using an extract which contains both carotenoids in the desired ratio, by adding an additional amount of one of the carotenoids to an extract comprising both carotenoids so that the desired ratio is obtained or by mixing the two separate carotenoids (each obtained by any of the methods mentioned or described above and below) to reach the desired ratio between them.

One of the novel features of the inventive composition is that while possessing the above noted properties, the combination of phytoene and phytofluene is essentially devoid of any color (but for a slight yellowish hew, hardly visible, of the phytofluene). The fact that the composition is essentially colorless ensures that these carotenoids will not have any effect on the aesthetic properties of the preparation comprising them. In addition, the lack of absorbance of light in the visible range (which is a manifestation of the fact that they are essentially colorless) renders them stable to degradation under visible light.

The preparation may, in accordance with the above noted preferred embodiment, be used as a topical cosmetic or pharmaceutical preparation in order to protect the skin from environmental hazards such as those described above including UV (UVA and/or UVB) irradiation or damages which can be effected by a variety of oxidative agents. A topical composition of the invention may be in the form of a gel, an oil-in-water or water-in-oil emulsion, a salve or ointment, etc.

In accordance with another embodiment, the composition may be used as an additive in food preparations, e.g. serving as a preservative to protect against oxidation of the various food ingredients, e.g. of oils or fats.

At times, the composition of the invention may comprise additional components which do not substantively change the basic characteristics of the composition. One example of such a component is zetacarotene (7,8,7',8' - tetrahydro- γ , γ -carotene).

The composition of the invention may obviously, depending on its use, comprise also other ingredients, cosmetical or pharmaceutical acceptable carriers, preservatives, other antioxidants, various pharmaceutically or cosmetically active ingredients such as a topically acting drugs, etc.

In accordance with one preferred embodiment, the composition comprises also a hydrophobic carrier, which may be selected from oils typically used in the cosmetic, pharmaceutical or food industry, such as vegetable, mineral or synthetic oils.

The phytoene and phytofluene may be obtained from a variety of sources. Typically, they may be obtained from organisms that produce carotenoids, such as a variety of plants, various algae, and particularly micro algae *Dunaliella sp*, being a specific example. Very low amounts of phytoene and phytofluene have been produced by β -carotenoid producing organisms. To obtain even such low amounts of the carotenoids, the organisms were grown under dim light conditions. In accordance with the invention, a method is provided which enables to yield substantive amounts of at least one of phytoene or phytofluene from such carotenoid producing organisms. The term "substantive amounts" relates to an

carotenoid producing organisms. The term "substantive amounts" relates to an amount of phytoene or an amount of phytofluene ranging from about 0.1 mg/l culture to about 30 mg/l culture typically between about 1 mg/l to about 20 mg/l. When the carotenoid producing organism is the algae *Dunaliella sp.*, the typical
5 amount of phytoene and phytofluene which may be produced is between about 1 mg/l to about 15 mg/l. In accordance with the method of the invention, the carotenoid producing organism, typically algae, may be grown under various kinds of conditions.

In accordance with one embodiment of this aspect of the invention, the
10 carotenoid producing organisms are grown outside, for example, in the case of *Dunaliella sp.* in outside pools in a growth medium comprising sea water and recycled salt water at a temperature of between about 10°C to about 40°C. The organisms may be grown under sunlight at an extent of between 80% shade to full sunlight. By a preferred embodiment, the carotenoid producing organisms are
15 grown under bright light ranging from about 40% shade to full sunlight.

By an additional embodiment, the carotenoid producing organisms are grown in a growth culture, typically, in a fermentor and typically at room temperature in which case the growth medium will typically comprise various salts and minerals in different combinations (see, for example, Example 1.1 below). In
20 accordance with this embodiment, the organisms are grown under broad spectrum light (light spanning over a substantive portion of visible light) of above about 10 W/m² and which may range from about 10 W/m² to about 200 W/m².

In accordance with the method of the invention, substantive amounts of phytoene and/or phytofluene are typically produced following several days of
25 growth of the organism (e.g. about 4 to 6 days).

The phytoene and phytofluene are obtained under conditions which favor the accumulation of phytoene and phytofluene in cells of the organisms, typically by growing a β -carotene-producing organism in the presence of carotenoid-biosynthesis inhibitors, which are inhibitors that can act in subsequent reaction
30 steps in the biochemical pathway of the production of carotenoids. One example of

inafter: "the 4-chloro inhibitor"). When the source of phytoene and phytofluene is the micro algae *Dunaliella*, the 4-chloro inhibitor is typically included in the growth medium in a concentration of about 0.1 μM (Ben-Amotz *et al.*, *Plant Physiol.* **86**:1286, 1988). Other inhibitors which may be used are, for example, J334,
5 (Ben-Amotz *et al. Supra*) Sandoz H 6706 (Kümmel, H.W. *et al.*, *Z. Naturforsch* **30c**, 333, 1975), di-phenyl-amine (DPA) (Foppen F.H., *Ann. Ist. Super. Sanita*, 439, 1969) and nicotine (Shaish *et al.*, *Plant Cell Physiol* **31**:689, (1990)).

In accordance with one embodiment of this aspect of the invention there is thus provided a method for preparing a composition comprising at least one of
10 phytoene or phytofluene, comprising the steps of:

- (i) incubating a carotenoid producing organism in a growth culture under broad spectrum light having an intensity of above about 10 W/m^2 in the presence of one or more carotenoid synthesis inhibitors;
- (ii) growing said carotenoid producing organisms until the level of at
15 least one of phytoene or phytofluene is between about 1 mg/l culture to about 50 mg/l culture; and
- (iii) separating said organisms from the culture.

In accordance with an additional embodiment of this aspect of the invention there is provided a method for preparing a composition comprising at least one of
20 phytoene or phytofluene, comprising the steps of:

- (i) Incubating a carotenoid producing organism in a growth culture under sunlight at an extent between about 80% shading to full sunlight in the presence of one or more carotenoid synthesis inhibitor;
- (ii) Growing said carotenoid producing organisms until the level of at
25 least one of phytoene or phytofluene is between about 1 mg/l culture to about 50 mg/l culture; and
- (iii) separating said organisms from the culture.

In accordance with this aspect of the invention, the phytoene and
30 phytofluene obtained in the grown organisms may be used without being separated

In accordance with this aspect of the invention, the phytoene and phytofluene obtained in the grown organisms may be used without being separated from the dry matter of the producing organism. However, in accordance with a preferred embodiment, following separation of the carotenoid producing organism
5 from the culture, a preparation is further prepared from the organisms comprising phytoene, phytofluene or both. Typically, the preparation is an extract which may be prepared from the carotenoid producing organisms by any of the extraction methods known in the art such as, for example, by ethanol:hexane extraction.

In accordance with the method of the invention, typically, both phytoene and
10 phytofluene are obtained from the carotenoid producing organism and the ratio of phytoene to phytofluene may vary according to the conditions under which the organism was grown and, when a preparation is prepared, the type of methods used for obtaining the preparation. At times, however, under certain conditions, growing the carotenoid producing organism in accordance with the method of the invention
15 may result in obtaining mostly one or only one of the two carotenoids.

In accordance with the invention it was also found that the carotenoids ratio may further be enhanced in favor of phytoene and phytofluene by the addition of active charcoal during carotenoid extraction. Charcoal may be added when an inhibitor is used as well as when the carotenoid producing organism is grown
20 without the inhibitor. The charcoal is later filtered out from the culture or preparation using any of the known centrifugation or filtration methods.

In addition to the above, in accordance with the invention the carotenoids in the composition of the invention may also be synthesized by any of the known chemical or biochemical methods or recombinant methods. Chemically, phytoene
25 can be synthesized, for example, from two geranylgeranyl pyrophosphates (C-20), in a reaction which may be mediated by phytoene synthase. The geranylgeranyl pyrophosphate can be obtained directly, by the conversion of mevalonic acid or by the condensation of pyrovate and glyceraldehyde-3-phosphate. Phytofluene can be synthesized by desaturation of phytoene, a reaction which may be mediated by
30 phytoene desaturase. Recombinant methods include, for example, the mutagenesis

of enzymes which are active downstream to phytofluene. Such synthesized phytoenes and phytofluenes will have activities which are substantively similar to the activities of the phytoene and phytofluene obtained from organisms that produce carotenoids as explained above.

5 The stability of the carotenoids in the composition of the invention can be tested by irradiating the carotenoid composition by light and/or exposing the composition to oxygenating agents. The effect of such treatments on the carotenoids in the compositions of the invention, is less than the effect of such treatments on other carotenoids such as, for example, β -carotene. In prior art
10 compositions containing an effective amount of carotenoids, such treatment typically result in degradation of the carotenoids, manifested for example by change in light absorbance, i.e. color. The anti-oxidative effect of the composition of the invention can be determined, for example, by determining the anti-oxidation protection effect on DNA (Salles *et al.*, *Anal. Biochem.*, 232:37, 1995).

15 The invention will now be illustrated further in the following description of the non-limiting specific embodiments.

BRIEF DESCRIPTION OF THE DRAWINGS:

Fig. 1 shows the UV absorbance spectra of a composition comprising
20 phytoene extracted from *Dunaliella sp.* grown with the 4-chloro inhibitor.

Fig. 2 shows the UV absorbance spectra of phytofluene extracted from *Dunaliella sp.* grown with the 4-chloro inhibitor.

EXAMPLES

25 I. MATERIALS AND METHODS

1. Growth mediums

The following growth mediums were used:

1.1 A growth medium comprising:

MgSO₄, 5 mM; CaCl₂, 0.2 mM; KH₂PO₄, 0.2 mM; FeCl₃ + Na₂EDTA, 2 μM+5 μM; MnCl₂, 7 μM; CuCl₂, 1 μM; ZnCl₂, 1 μM; CoCl₂, 1 μM; (NH₄)₆MO₇O₂₄, 1 μM; NaCl, 1M; NaHCO₃, 50 mM; KCl, 5 mM.

The growth medium is typically used for growth of carotenoid producing organisms inside under artificial conditions.

1.2 A growth medium comprising sea water and recycled salt water in an amount which results in 2M salinity. Such a medium is typically used while growing a carotenoid producing organism outside under sunlight.

The final pH of the growth mediums was 7.0 to 8.0.

10

2. Chemicals

β-Carotene biosynthesis inhibitor 4-chloro-5(methylamino)-2-(3-(trifluoromethyl)phenyl)-3(2H)-pyridazinone. The 4-chloro inhibitor was used for phytoene and phytofluene production. The range of concentration of the 4-chloro inhibitor was between about 0.07 μM to about 0.5 μM.

15

Untreated active charcoal was used to increase the ratio of phytoene and phytofluene to other carotenoids. The charcoal was filtered out from the extract at the final steps of preparation of the extract.

Various commercially used oils were used to dissolve the phytoene and phytofluene.

20

3. Growth of *Dunaliella* sp.

The algae *Dunaliella* sp was grown under one of the following conditions:

3.1 In the growth medium described in 1.1 above in a fermentor at room temperature under artificial lighting which altered in the range of between dim light (from about 1 W/m²) to bright light (above about 10 W/m²), typically under light of above 10 W/m² in the range of between 10 W/m² to about 200 W/m².

25

3.2 In outside pools at a temperature of between about 10°C to about 40°C under light conditions in the range of between 80% shade to full sunlight, typically between 40% shade to full sunlight.

5 4. Carotenoids extraction method

Phytoene and phytofluene were produced by inhibition of β -carotene synthesis in the algae *Dunaliella sp.* By the 4-chloro inhibitor at a concentration of 0.07 μ M to 1.0 μ M.

The algae were collected after four to six days of growth of the algae either
10 outdoors or indoors as described above and the cell pellet was extracted with ethanol:hexane (1:2) v/v. Ethanol was first added to the cell pellet at algae:ethanol ratio of at least 1:10. At this stage basic hydrolysis of ester bonds is performed by the addition of 0.5M NaOH with stirring for at least 30 min. Ethanol:hexane phase separation is achieved by the addition of NaCl at adequate quantity. The hexane
15 fraction was analyzed spectrophotometrically.

Hexane was dried under vacuum and the carotenoids were re-dissolved in hexane or oil.

5. Spectrophotometric analysis

20 Absorption spectra of the phytoene and phytofluene extract was determined using Hewlett Packard 8452A Diode Array Spectrophotometer.

6. Stability of phytoene and phytofluene under aerobic conditions in UV lights and various incubation temperatures

25

Stability of the carotenoids in the various temperatures was determined by long-term incubation of phytoene and phytofluene in several types of oils and solvents at 4°C, 23°C, 30°C and 60°C. The amount of phytoene and phytofluene left was determined spectrophotometrically. The percent of phytoene and
30 phytofluene left was calculated as follows:

$$\frac{\text{Phytoene and phytofluene after treatment (mg/ml)} * 100}{\text{Phytoene and phytofluene before treatment (mg/ml)}} = \% \text{ PH and PF left}$$

- 5 Stability under UV lights (254 nm, 365 nm, 254 + 365 nm), for 30 mins. at 300/310 $\mu\text{W}/\text{cm}^2$ was determined.

7. **Stability of phytoene and phytofluene under aerobic conditions in visible light**

10

Stability of phytoene and phytofluene under visible light was measured under sunlight (full sun summer day in Rehovot, Israel, at noon) filtered through "hot mirror" (Andover Corporation, Salem, NH, transmittance 400-650nm) filter. Samples were dissolved in ethanol, exposed to light for 30-150 min and then
15 extracted in hexane after basic hydrolysis (as in extraction method above). Phytoene, phytofluene and beta-carotene amounts were measured spectrophotometrically, and percentage lefts were calculated as in section 5 above.

8. **Anti-free radical activity**

20

8.a Quenching of hydroxyl radicals:

The ability of the compound to quench the activity of hydroxyl radicals ($^{\circ}\text{OH}$) was measured by Electron Paramagnetic Resonance (EPR), by comparing the signal intensity with and without the compound. Hydroxyl radicals were generated by decomposition of hydrogen peroxide by iron
25 $[(\text{FeSO}_4) \text{ Fenton reaction}]$. The $^{\circ}\text{OH}$ is trapped by the DMPO to provide a DMPO-OH adduct, which presents a characteristic signal in EPR. The compound, composed of phytoene and phytofluene at a ratio of 6.66:1 respectively, was added to the system in three final dilutions of 1/20, 1/50 and 1/200.

30

Since various substances can reduce the signal, not by trapping of the $^{\circ}\text{OH}$ but by addition to the DMPO-OH adduct, it is necessary to also evaluate the signal when the product is added after the Fenton reaction

(post-addition). The trapping capacity of the hydroxyl radical is evaluated theoretically by the difference between the value of the EPR signal corresponding to pre-addition of the product and the value of the signal resulting from post-addition.

8.b DNA protection by anti-oxidation effects:

DNA protection by the anti-oxidation effect of the compound was examined based on the method of Salles *et al.* (*Supra*). In the test the protective effect of the compound against Reactive Oxygen Species (ROS) generated oxidative DNA damage was measured, by quantifying the inhibition of DNA lesions formation using plasmid DNA target.

II. RESULTS

Example 1 Phytoene and phytofluene production

Phytoene and phytofluene were produced from the algae *Dunaliella sp.* by growing the algae in the presence of 0.1-0.5 μg of the β -carotene biosynthesis 4-chloro inhibitor as described above. The cultures were then extracted and the extract was dried by evaporation and re-dissolved in hexane or oil. No residues of the 4-chloro inhibitor were detected in the extracted carotenoids.

As seen in Fig. 1, the main absorbance peak of phytoene was found to be at 286 nm (UVB) and as seen in Fig. 2, the main absorbance peak of phytofluene was found to be at 348 nm (UVA). As expected, the main absorbance peak for β -carotene was at 450 nm. β -Carotene was reduced due to the synthesis inhibition while phytoene and phytofluene increased.

Example 2 Example for the stability of the extracted phytoene and phytofluene in UV lights

The stability of the phytoene and phytofluene extracts obtained as explained in Example 1 above was determined by spectrophotometric analysis. Phytoene and

phytofluene were dissolved in hexane or oil. The absorbance of the preparation is determined in 220-600 nm and the amount of each of the above carotenoids in each sample is calculated to determine the percent of the carotenoids measured after irradiation of the extract divided by the amount of the same carotenoid which was measured in the extract before exposing it to irradiation (see Materials and Methods 5 above). The stability of the phytoene and phytofluene was measured by the exposure of the extract both to UVA light (365 nm) and UVB light (254 nm) irradiation as well as to a combination of UVA and UVB irradiation.

As seen in Table 1 below, phytoene and phytofluene dissolved in oil or 10 hexane were very stable after exposure to UVA, UVB or UVA+B irradiation.

Table 1

15 **PH and PF stability in UV irradiation (30 minutes exposure)**

	Darkness	254 nm	365 nm	254nm+365nm
	PH and PF (% left) ¹	PH and PF (% left)	PH and PF (% left)	PH and PF (% left)
PH left ¹ in hexane	100	92.2	98.7	92.4
PF left ¹ in hexane	100	92.6	101.5	99.0
PH left ¹ in oil	100	100	100	101
PF left ¹ in oil	100	90	88	98

¹ Percent of PH or PF left is the amount of material (mg/ml) measured after irradiation divided by the amount of material measured before treatment multiply by 100.

20

The stability of phytoene and phytofluene under different temperature conditions (4°C, 23°C, 30°C and 60°C) in hexane or in other various commercially used oils was also measured.

The measurements were carried out over a period of four months.

Phytoene and phytofluene were found to be stable in the entire measured temperature range with no substantial effect of the kind of oil in which the carotenoids were dissolved.

5

Example 3 Stability of the extracted phytoene and phytofluene in visible light

Stability of phytoene and phytofluene was compared to that of beta-carotene stability (as explained in materials and methods) under visible light. The absorbance of the preparation is determined in 220-600 nm wavelength and the amount of each of the above carotenoids in each sample is calculated to determine the percent of the carotenoids measured after irradiation of the extract compared to measurements prior to irradiant (see example 2 above).

15

As shown in table 2, exposure to visible high intensity light cause significant higher degradation of beta-carotene compared to phytoene and phytofluene.

Table 2

PH and PF stability in visible light (15-150 min exposure)

Time of exposure (min)	Phytoene (% left)	Phytofluene (% left)	Beta-carotene (% left)
0	100	100	100
30	54.46	66.13	25.00
90	20.33	50.42	11.36
120	15.70	41.33	9.47
150	13.64	33.06	5.68

20

Example 4 Anti free radical activity

4.a Quenching of hydroxyl radicals:

25

Quenching of hydroxyl radicals was measured by EPR as explained in materials and methods. The results obtained (Table 3) indicate a

dose-dependent trapping activity of the hydroxyl radical by the product.

Table 3

Product final concentration (mg/ml)	Intensity of EPR signal (arbitrary units)	Intensity of the EPR signal after post-addition (arbitrary units)	% of hydroxyl radicals trapping
0	$2.82 \times 10^6 \pm 0.07$		
0.001	$0.34 \times 10^6 \pm 0.04$	$2.19 \times 10^6 \pm 0.13$	66
0.0004	$0.58 \times 10^6 \pm 0.02$	$1.94 \times 10^6 \pm 0.05$	48
0.0001	$1.26 \times 10^6 \pm 0.10$	$2.19 \times 10^6 \pm 0.13$	24

5

The product was composed of phytoene and phytofluene at a ratio of 6.66:1 respectively at an initial concentration of phytoene 0.02 mg/ml and phytofluene 0.003 mg/ml.

The average value of each result corresponds to three measurements.

10

4.b Anti oxidation effect:

DNA protection by the anti oxidation effect of the phytoene and phytofluene extracts was determined based on the Salles method (*Supra*) as explained in Materials and Methods 7.b above.

As seen in Table 4 below, phytoene and phytofluene obtained as described
 15 above were capable of protecting against hydroxyl radicals.

Table 4
PH and PF anti oxidation activity

Compound	Concentration μg/ml	% of inhibition in the presence of ROS	% of non specific inhibition	% of specific inhibition	Concentration giving 50% of activity
Phytoene and phytofluene	140	86	24	62	11.2 μg/ml
	14	74	19	55	
	1.4	6	7	1	
	0.14	1	1	0	
Positive control	1000	88	9	79	80 μg/ml
	100	69	11	58	
	10	5	0	5	
	1	4	0	4	

5 Example 5 Protective effects on DNA by phytoene and phytofluene

Genotoxic effect

The genotoxicity is measured as the induction of DNA repair synthesis activity and is expressed by the ratio (R): while $R \leq 2$ indicate no toxic effect

$$10 \quad R = \frac{\text{Relative Light Unit of the sample}}{\text{Relative Light Unit of the solvent alone}}$$

As seen in Table 5 below, phytoene and phytofluene are not genotoxic.

15 The R parameter is significantly less than 2 (protective effect) and much smaller in comparison with the positive control, which is a known genotoxic compound.

Table 5
Genotoxic effect of phytoene and phytofluene

Compound	Concentration ($\mu\text{g/ml}$)	Genotoxicity (Ratio)*
Phytoene and phytofluene	140	0.92
	14	1.11
	1.4	1.18
	0.14	0.05
MMS-(positive control)	10 mM	6.26
	2 mM	2.44

* See above

5 Example 6

The following are examples of compositions that may be used in accordance with the invention:

A- *Emulsified gel of O/W* (topical route):

10	- Carbopol 981 (marketed by Goodrich).....	0.6	g
	- Ethyl alcohol.....	15	g
	- Volatile silicone oil.....	3	g
	- Purcellin oil.....	7	g
	- Preserving agent.....	0.3	g
15	- Perfume.....	0.4	g
	- Triethanolamine.....	0.2	g
	- Phytoene	0.01	g
	- Phytofluene.....	0.001	g
	- Demineralized water.qs	100	g

20

B - *Anhydrous gel* (topical route):

	- Propylene glycol.....	25	g
	- Hydroxyethyl cellulose.....	0.8	g
25	- Polyethylene glycol	12	g
	- Phytoene.....	1	g
	- Phytofluene.....	1	g

- Absolute ethanol.qs..... 100 g

C- Emulsion of O/W type (topical route):

- Liquid paraffin..... 6 g
- 5 - Liquid lanolin..... 3 g
- Arlacel 165 (marketed by Atlas) 6 g
- Tween 60 (marketed by Atlas) 2 g
- Cetyl alcohol..... 1.2 g
- Stearic acid..... 2.5 g
- 10 - Volatile silicone oil..... 10 g
- Triethanolamine..... 0.1 g
- Preserving agent..... 0.3 g
- Antioxidants..... 0.3 g
- Phytoene 0.3 g
- 15 - Phytofluene 0.2 g
- Demineralized water.qs 100 g

D - Cream containing liposomes (topical route):

- Sunflower oil 35 g
- 20 - Cetyl alcohol..... 4 g
- B-sitosterol..... 4 g
- Dicetyl phosphate 0.5 g
- Preserving agent..... 0.3 g
- Perfume..... 0.6 g
- 25 - Carbopol 981 (marketed by Goodrich)..... 0.2 g
- Triethanolamine..... 0.2 g
- Sphingosine..... 0.05 g
- Phytoene 0.00001 g
- Phytofluene 0.000001 g
- 30 - Demineralized water.qs 100 g

E - Per os composition:

- 19 -

-	Talc	5 g
-	Aerosil 200.....	5 g
-	Stearate de Zn	5 g
-	Phytoene	0.0000015 g
5 -	Phytofluene.....	0.000001 g
-	Lactose qs	400 g

F – *Emulsion W.O* (topical route):

-	Protegin (marketed by Goldschmidt)	19 g
10 -	Glycerine.....	3 g
-	Vaseline oil	8 g
-	Phytofluene.....	0.5 g
-	Phytoene	0.5 g
-	Sulfate de Mg.....	0.5 g
15 -	Perfume.....	0.8 g
-	Preserving agent.....	0.2 g
-	Water.qs	100 g

CLAIMS:

1. A composition comprising, an amount of phytoene and an amount of phytofluene, effective in combination in the prevention of damage resulting from oxidation or from exposure to UV light.
2. A composition according to Claim 1, being a topical skin preparation for protecting the skin against environmental hazards and comprising phytoene and phytofluene in an effective amount such that, in combination, these carotenoids exert an oxidation-protecting and UV-protecting effect on the skin.
3. A composition according to Claim 2, being a cosmetic or pharmaceutical composition.
4. A composition according to Claim 1, being a food additive.
5. A composition according to any of the previous claims being essentially colorless.
6. A composition according to any of the previous claims, wherein the weight ratio between the phytoene and phytofluene in the composition is in the range of about 200:1 to about 1:200, respectively.
7. A composition in accordance with Claim 6, wherein the weight ratio between phytoene and phytofluene is about 50:1 to about 1:50, respectively.
8. A composition in accordance with Claim 7, wherein the weight ratio between phytoene and phytofluene in the composition is in the range of between about 10:1 to about 1:10, respectively.
9. A composition according to any of the previous claims, comprising a hydrophobic carrier.
10. A composition according to any one of the previous claims further comprising zetacarotene.
11. A method for preparing a composition comprising at least one of phytoene or phytofluene, comprising the steps of:

- (i) incubating a carotenoid producing organism in a growth culture under broad spectrum light having an intensity of above about 10 W/m² in the presence of one or more carotenoid synthesis inhibitors;
- (ii) growing said carotenoid producing organisms until the level of at least one of phytoene or phytofluene is between about 1 mg/l culture to about 50 mg/l culture; and
- (iii) separating said organisms from the culture.

12. A method according to Claim 11, wherein said light has an intensity in the range of between about 20 W/m² to about 200 W/m².

10 13. A method for preparing a composition comprising at least one of phytoene or phytofluene, comprising the steps of:

- (i) Incubating a carotenoid producing organism in a growth culture under sunlight at an extent between about 80% shading to full sunlight in the presence of one or more carotenoid synthesis inhibitors;
- (ii) growing said carotenoid producing organisms until the level of at least one of phytoene or phytofluene is between about 1 mg/l culture to about 50 mg/l culture; and
- (iii) separating said organisms from the culture.

20 14. A method according to Claim 13, wherein the extent of sunlight is between about 80% shading to full sunlight.

15. A method according to Claim 13, wherein said organisms are grown under full sunlight.

16. A method according to any of Claims 11-15, comprising the additional step of:

- (iv) preparing a preparation comprising phytoene, phytofluene or both from said separated organism.

17. A method according to Claim 16, wherein said preparation is an extract.

18. A method according to Claim 17, wherein said extract is an ethanol:hexane extract.

19. A method according to any of Claims 11-18, wherein said carotenoid producing organism is a micro algae.
20. A method according to Claim 19, wherein said micro algae is *Dunaliella* sp.
21. A method according to any of Claims 11-20, wherein said carotenoid
5 synthesis inhibitor is 4-chloro-5(methylamino)-2-(3-(trifluoromethyl)phenyl)-
3(2H)-pyridazinone.
22. A method according to any of Claims 16-21, wherein said extract comprises
no traces of said inhibitor.
23. A method in accordance with any of Claims 16-22, further comprising the
10 addition of charcoal to said carotenoid preparation during its preparation.
24. A composition in accordance with any of Claims 1-10, wherein said
phytoene and phytofluene are prepared by chemical synthesis.
25. A composition in accordance with any of Claims 1-10, wherein said
phytoene and phytofluene are prepared by recombinant methods.

1 / 1




FIG.1




FIG.2

1 / 1




FIG.1





FIG.2



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A23L 1/303, A61K 7/00, 7/06, 7/42, 7/48	A3	(11) International Publication Number: WO 00/13654 (43) International Publication Date: 16 March 2000 (16.03.00)
(21) International Application Number: PCT/IL99/00478 (22) International Filing Date: 5 September 1999 (05.09.99) (30) Priority Data: 126076 4 September 1998 (04.09.98) IL (71) Applicant (for all designated States except US): I.B.R. ISRAELI BIOTECHNOLOGY RESEARCH, LTD. [IL/IL]; 85 Derech Namir, Apt. 23, 62301 Tel Aviv (IL). (72) Inventors; and (75) Inventors/Applicants (for US only): SOUDANT, Etienne [FR/FR]; 14, rue Leon Bernard, F-94260 Fresnes (FR). BEZALEL, Lea [IL/IL]; Derech Mezada 15, 84359 Beer Sheva (IL). SCHICKLER, Hedva [IL/IL]; 19 Hamagen Street, 76804 Mazkeret Batya (IL). PALTIEL, Judith [IL/IL]; 12/26 Parshani Street, 76000 Rehovot (IL). BEN-AMOTZ, Ami [IL/IL]; 21 Hatomer Street, 56542 Savion (IL). SHAISH, Aviv [IL/IL]; 37 Achva Street, 79800 Shikmim (IL). PERRY, Inon [IL/IL]; 85 Derech Namir, Apt 23, 62301 Tel Aviv (IL). (74) Agent: REINHOLD COHN AND PARTNERS; P.O.Box 4060, 61040 Tel Aviv (IL).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 24 August 2000 (24.08.00)
(54) Title: A CAROTENOID COMPOSITION CONTAINING PHYTOENE AND PHYTOFLUENE (57) Abstract <p>A preparation comprising the carotenoids phytoene and phytofluene in an amount which, in combination, is effective in preventing various kinds of damage resulting from oxidation and exposure to UV light is provided. The preparation is essentially colorless and may be used as a topical cosmetic or pharmaceutical preparation as well as an additive in food preparations. In addition, a method for producing substantive amounts of phytoene and phytofluene from carotenoid producing organisms is described as well.</p> <div style="text-align: center;">  <p>286</p> <p>2.5</p> <p>0.0</p> <p>300</p> <p>WAVELENGTH (nm)</p> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

Int. l. Application No

PCT/IL 99/00478

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A23L1/303 A61K7/00 A61K7/06 A61K7/42 A61K7/48		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 A23L A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TONUCCI, L.H. ET AL.: "Carotenoid Content of Thermally Processed Tomato-Based Food Products" J. AGRIC. FOOD CHEM., vol. 43, no. 3, 1995, pages 579-586, XP002124250	1,4,10, 24,25
Y	page 584, column 1, line 31 - line 39; tables 2,3 <div style="text-align: center; margin-top: 10px;"> --- -/-- </div>	1-10
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
* Special categories of cited documents :		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"G" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search	Date of mailing of the international search report	
14 Apr11 2000	31.05.00	
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016		Authorized officer <div style="text-align: center; margin-top: 10px;">Minas, S</div>

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 99/00478

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHOO, Y.M.: "Production of palm oil carotenoid concentrate and its potential application in nutrition" STN INTERNATIONAL; FILE CHEMICAL ABSTRACTS, AN119:124982, 1992, XP002124251	1,4,10, 24,25
Y	abstract & LIPID-SOLUBLE ANTIOXID. EDITOR(S): ONG, AUGUSTINE S.H.; PACKER, LESTER. PUBLISHER: BIRKHÄUSER, BASEL, SWITZ, 1992, pages 243-254,	1-10
X	WO 96 39869 A (MARS INC) 19 December 1996 (1996-12-19)	1,4,9, 10,24,25
Y	page 4, line 36 -page 5, line 5; claims 1,3,5,7,9; examples 6,7	1-10
Y	FR 2 759 904 A (REZA ALMA HENY) 28 August 1998 (1998-08-28) claims 1,19,24,25; example 3	1-3,9,10
Y	PATENT ABSTRACTS OF JAPAN vol. 005, no. 168 (C-077), 27 October 1981 (1981-10-27) & JP 56 099405 A (NISSHIN OIL MILLS LTD:THE), 10 August 1981 (1981-08-10) abstract	1-3,9,10
A	WO 93 04598 A (BETATENE LTD) 18 March 1993 (1993-03-18) claims 1,10,19	1-10
X	EP 0 769 551 A (KIRIN BREWERY) 23 April 1997 (1997-04-23) abstract	1
X	US 5 545 816 A (AUSICH RODNEY L ET AL) 13 August 1996 (1996-08-13) abstract	1
X	BEN-AMOTZ, A. ET AL.: "The biotechnology of cultivating Dunaliella for production of beta-carotene rich algae" BIORESOURCE TECHNOLOGY, vol. 38, 1991, pages 233-236, XP000901780	11,13, 19,20
Y	the whole document	11-21
Y	BEN-AMOTZ, A. ET AL.: "Massive accumulation of phytoene induced by norflurazon in Dunaliella bardawil (chlorophyceae) prevents recovery from photoinhibition" JOURNAL OF PHYCOLOGY, vol. 23, no. 1, March 1987 (1987-03), pages 176-181, XP000901498 the whole document	11-21

INTERNATIONAL SEARCH REPORT

PCT/IL 99/00478

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	VAISBERG, A. J. AND SCHIFF, J. A.: "Events surrounding the early development of Euglena chloroplast" PLANT PHYSIOLOGY, vol. 57, 1976, pages 260-269, XP000901340 the whole document	11-15, 21
Y	US 4 199 895 A (AVRON MORDHAY ET AL) 29 April 1980 (1980-04-29) claims 1, 3, 8, 10	13-21
Y	US 5 310 554 A (HAIGH W GEOFFREY) 10 May 1994 (1994-05-10) claims 1-7; example 1	11, 16-21
Y	FR 2 620 131 A (COMMISSARIAT ENERGIE ATOMIQUE) 10 March 1989 (1989-03-10) page 1, line 31 - line 32 page 5, line 32 - line 35; claims 1, 13	11-15, 19-21
Y	FR 2 678 946 A (OVI) 15 January 1993 (1993-01-15) page 5, line 6 - line 15 page 9, line 30 - line 36 page 12, line 30; claims 1, 12	11-15, 19-21
A	DD 246 565 A (PAEDAGOGISCHE HOCHSCHULE WOLFG) 10 June 1987 (1987-06-10) page 2, line 35; claim 3	11-23

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IL 99/00478

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-10, 24, 25

A composition comprising phytoene and phytofluene.

2. Claims: 11-23

A method for producing and extracting phytoene or phythofluene.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inten. Appl. No.

PCT/IL 99/00478

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9639869 A	19-12-1996	US 5643623 A AU 5880696 A EP 0831726 A JP 11506925 T PL 324969 A US 5834044 A	01-07-1997 30-12-1996 01-04-1998 22-06-1999 22-06-1998 10-11-1998
FR 2759904 A	28-08-1998	CA 2252633 A EP 0917456 A WO 9837864 A	03-09-1998 26-05-1999 03-09-1998
JP 56099405 A	10-08-1981	JP 1400944 C JP 62006682 B	28-09-1987 13-02-1987
WO 9304598 A	18-03-1993	AU 654876 B EP 0602137 A JP 7501211 T	24-11-1994 22-06-1994 09-02-1995
EP 0769551 A	23-04-1997	JP 8242861 A AU 685354 B AU 4889996 A NO 964754 A WO 9628545 A	24-09-1996 15-01-1998 02-10-1996 08-01-1997 19-09-1996
US 5545816 A	13-08-1996	US 5618988 A CA 2055447 A EP 0471056 A JP 5504686 T WO 9113078 A US 5530188 A US 5530189 A US 5684238 A US 5656472 A	08-04-1997 03-09-1991 19-02-1992 22-07-1993 05-09-1991 25-06-1996 25-06-1996 04-11-1997 12-08-1997
US 4199895 A	29-04-1980	IL 54881 A US 4511 P	29-06-1981 18-03-1980
US 5310554 A	10-05-1994	AU 681868 B AU 5453094 A CA 2147825 A EP 0666845 A JP 9500864 T WO 9410140 A	11-09-1997 24-05-1994 11-05-1994 16-08-1995 28-01-1997 11-05-1994
FR 2620131 A	10-03-1989	AU 606458 B AU 2322288 A EP 0329754 A WO 8901977 A IL 87576 A JP 2501189 T NO 891745 A	07-02-1991 31-03-1989 30-08-1989 09-03-1989 10-06-1993 26-04-1990 26-04-1989
FR 2678946 A	15-01-1993	NONE	
DD 246565 A	10-06-1987	NONE	